



Tips for helping your child through a traumatic event

When traumatic events take place, children do not have the ability to communicate worries or concerns through words, they have not yet developed the vocabulary. Teens have the vocabulary, but may lack the maturity and life experience to properly identify the feelings they are having.

A traumatic event is frightening. It can be even more frightening and unsettling for children. Children experience the same event and are exposed to the same photographs and stories in print and television media.

Children are like sponges, taking in the world. They are very aware when the adults around them are having a strong emotional reaction. They need to have their feelings validated. It is important to remember that a child not talking about an event does not mean that the event is forgotten or has no importance. Children and teens are more likely to express their reactions through their actions rather than words. Be attentive and take special care through this time of chaos and instability.

The children in your life need your reassurance that they are safe with you. It is important to let them know that it is healthy to express emotions and to let them out. During this time, children will likely be highly sensitive and attentive to the adult responses surrounding them.

The children in your care will need reassurance that their safety and security will be taken care of by the adults in their lives. They will need encouragement to express their concerns. An extra hug, talk time or five more minutes before bedtime are some examples of what you can do to encourage and reassure them. Let them tell you what they need by paying attention to what they are doing or asking for.

Potential reactions that you might see in your child

- Irritability, agitation, confusion
- Withdrawn, denial, sadness
- Sleep disturbance, nightmares
- Increased anxiety, intense fear
- Changes in appetite
- Distress due to work or travel related parental absence
- Testing house rules-dress, curfew, chores
- Quietness, less communicative
- Feelings of helplessness or anger
- More prone to accidents/clumsiness
- Verbal outbursts
- Poor concentration
- Return/increase of bed-wetting or other regressive behavior

The reactions are usually temporary and will often subside in three to six weeks. In the meantime, these reactions can be uncomfortable, impacting concentration, productivity and even disrupting sleep patterns.

What you can do

- Talk to your child about the situation
- Encourage your child to talk
- Say it is OK to be afraid
- Tell your child about your plans each day
- Be attentive to media coverage adding to you and your child's anxiety/stress level
- Put order in your day as best you can
- Explain, talk and listen frequently
- Stay together to show you won't go away
- Hold and hug
- Listen
- Let them know that you were afraid too
- Reassure them that they're in a safe place
- As long as it can be done safely, let your child help clean up

Bedtime may be difficult. Your child may be afraid of the dark, not want to sleep away from you and/or may have nightmares or begin to wet the bed again.

What you can do at bedtime

- Explain why it gets dark
- Praise the child for good things
- Leave the light on and the door ajar
- Read to your child
- Talk about dreams
- Don't yell
- Agree on a time for your child to go to bed
- Tell a story about something good that happened that day

It is important to help your child externalize the feelings. Keeping reactions inside will only serve to have them go into hiding and come back when least expected. Also, speak with other parents, caregivers, your child's teacher or school principal to discuss other ideas and suggestions, support and resources.

Take care of yourself

Be aware of your own stress reactions in order to take care of yourself. Your emotional health is an essential element to providing appropriate support to your children. Some final tips for you;

- If you wake up in the middle of the night have a drink of water, read a little if you can't get back to sleep and remember to take some deep breaths.
- If you have a flashback, take some deep breaths, breathing in through your nose, out

through your mouth to the count of ten, and take notice of your current environment.

- Get rest, keep a normal schedule and eat properly.
- Try to exercise where possible, drink lots of water and avoid alcohol.
- Connect with and get support from the people and resources that make a difference in your life.